



575 South 70<sup>th</sup> Street  
Suite 200  
Saint Elizabeth Medical Plaza  
Lincoln, Nebraska 68510-2471

Toll-Free (888) 488-6667  
(402) 488-3322  
Fax (402) 488-1172  
[www.nebraskaortho.com](http://www.nebraskaortho.com)

Patrick E. Clare, M.D. - Emeritus  
Ronald O. Schwab, M.D. - Emeritus  
Donald J. Walla, M.D.  
Thomas M. Heiser, M.D.  
Daniel R. Ripa, M.D.

Robert W. Dugas, M.D.  
David J. Clare, M.D.  
James W. Gallentine, M.D.  
Steve J. Volin, M.D.

Justin D. Harris, M.D.  
Scott A. Swanson, M.D.  
Daniel B. Cullan II, M.D.  
Aaron M. Bott, M.D.

Joseph P. Mulka, M.D., Ph.D.  
Sukchan Lee, M.D.  
Dane C. Todd, M.D.  
David P. Heiser, M.D. - consulting

## MPFL Reconstruction Instructions

Dane Todd, MD

1. **Elevate and Ice** the knee for 72 hours after surgery. The knee should be at or above the level of your heart. Do not place any pillows behind the knee. Pillows should be behind the foot.
2. **Pain Control** – You will be sent home with a prescription for pain medication. Take it according to the instructions on the prescription. Pain can cause constipation, so be sure to drink plenty of fluids and use an over the counter stool softener (i.e. Miralax) to help prevent this. You may take Tylenol (acetaminophen) instead of the prescribed medication. However, make sure you do not take more than 3,000 milligrams (3 grams) of Tylenol (acetaminophen) in any 24 hour period. Wean off the pain medications as soon as possible after surgery.
3. **Dressings** – Keep your dressings on for 3 days after surgery. Then you may remove the dressings to shower. Please re-apply new gauze dressings over the wound. Do not use any ointment, creams, or lotions on the incision. Leave the steri-strips in place. They will fall off on their own.
4. **Showering** – You may shower 3 days after surgery. Wash your incision gently with soap and water; do not scrub the incision. Pat the wound dry and apply new gauze. Do NOT take a bath or submerge your wound under water (i.e. pools, hot tub).
5. **Weight Bearing** – Touch Toe Weight Bearing for 1 week. After One week weight bearing as tolerated WITH crutches and brace locked in extension (for a total of 6 weeks).
6. **Range of Motion** – Keep in full extension for the first 2 weeks with no range of motion.
7. **Brace** – Wear at all times except when showering for the first 2 weeks.
8. **Physical Therapy** – Begin Physical therapy 2 weeks after surgery. Take the Physical Therapy prescription you were given to a therapist near your home to begin therapy.
9. **Driving** – You may drive once you are completely off all pain medications and feel comfortable doing so. Make sure to leave yourself extra space and time to stop and avoid any sudden movements.